



MATRIX EXPONENTIAL–BASED METHODS FOR SOLVING SYSTEMS OF DIFFERENTIAL EQUATIONS

Dr. Ajay Kumar Mishra

Associate Professor

Mathematics

Hierank Business School, Noida

ABSTRACT

Differential equations are systems of equations of central mathematical interest in the modeling of dynamics in science and engineering, economics, and applied mathematics. The number of available analytical and numerical methods has been quite high, but the procedures based on matrix exponential are a formidable and unified approach to the problem of linear and semi-linear systems of ordinary differential equations. These techniques provide precise solution approximations of linear problems and form the theoretical basis of the current exponential integrators applied to stiff and large-scale problems. This paper is a thorough and in-depth research on the techniques of solving exponential matrices, their mathematical basis, methods of finding the analytic solution, numerical estimates, and their applications. Mathematical properties of matrix exponentials are worked out, and then exponential time differencing, Krylov subspace methods, and formulations of low rank are discussed. To illustrate the depth of the theory as well as its practical applicability, illustrative examples, theorems, and proofs are presented. The paper shall be self-sustaining and fit for peer-reviewed publication and presentation in academic databases like ResearchGate.

Keywords: *Matrix Exponential, Systems of Differential Equations, Exponential Integrators, Krylov Subspace Methods, Numerical Analysis.*

1. INTRODUCTION

Ordinary differential equations (ODEs) are systems of mathematical equations that occur naturally in mathematical models of time-depending processes in a very broad variety of fields, such as physics and engineering, biology and economics, as well as control theory. These systems can be written in a compact form in matrices-vector in a large number of real world cases, such that one relies on linear algebraic tools to take a central role in the theoretical and computational strategy of solution.

The classical methods in solving systems of differential equations are eigenvalue eigenvector methods, Laplace transforms and numerical time stepping methods like Euler and Runge Kutta methods. Although these methods are useful in low-dimensional or non-stiff cases, they are usually insufficient in large, stiff, and highly structured systems. Specifically, classical numerical methods can need step sizes which are prohibitively small in order to be stable and accurate.

The methods based on matrix exponential are an elegant alternative and mathematically rigorous. The methods can give precise approximations of solutions to linear systems of differential equations by applying the same scalar operation, exponential, to a matrix argument. The matrix exponential is the operator that captures the full dynamical dynamics of a linear system and is the basic operator which dictates time dynamics. This renders it an intuitive and strong device of analytical studies as well as of numerical algorithms.

This paper is intended to give a rigorous and detailed treatment, mathematically, of methods of solving systems of differential equations based on the use of matrix exponentials. The paper lays out the theoretical basis of matrix exponentials, develops a representation of solutions to both non-homogeneous and homogeneous systems, and analyzes more sophisticated numerical techniques of computation efficiency. Clarity, rigor, and completeness are given importance and thus the work can be easily published and disseminated in the peer-reviewed journals and other academic platforms.

2. LITERATURE REVIEW

Baleanu et al. (2018) investigated a Chebyshev spectral method based on an operational matrix for solving fractional differential equations involving a non-singular Mittag-Leffler kernel and developed a robust numerical framework for handling such complex equations. The study demonstrated that the use of spectral methods significantly improved the accuracy and convergence rate of solutions while minimizing computational effort. It also highlighted that the proposed technique was particularly suitable for modeling real-world phenomena described by fractional dynamics, thereby contributing to advancements in applied mathematics, physics, and engineering.

Bloomfield (2018) explored the application of the R programming language in numerical analysis for science and engineering and demonstrated its versatility as a computational tool. The work provided detailed insights into how R could be used to solve differential equations, perform simulations, and analyze complex datasets. It emphasized that R's flexibility, wide range of libraries, and user-friendly

environment made it highly effective for mathematical modeling, enabling researchers and students to implement numerical methods efficiently and accurately.

Margossian (2019) reviewed the concept of automatic differentiation and its efficient implementation, focusing on its role in improving the precision and performance of computational methods. The study explained that automatic differentiation allowed exact derivative computation without the approximation errors associated with numerical differentiation. It also discussed different implementation strategies, such as forward and reverse modes, and highlighted their importance in large-scale optimization, machine learning, and scientific computing, where accurate gradient evaluation is critical.

Mena et al. (2018) examined numerical low-rank approximation techniques for solving matrix differential equations and proposed efficient computational methods for handling high-dimensional systems. The study showed that low-rank approaches significantly reduced memory requirements and computational complexity while preserving solution accuracy. It further emphasized that these methods were highly suitable for large-scale problems in engineering, physics, and applied mathematics, where traditional numerical techniques often become inefficient, thereby contributing to the development of scalable and efficient solution strategies.

3. MATHEMATICAL PRELIMINARIES

This section will focus on the fundamental mathematical knowledge and findings needed to develop the matrix exponential based methods of solving systems of differential equations. This conversation is about constant coefficient linear systems and defines the properties of matrix exponential which will form the basis of the rest of this paper.

3.1 Linear Systems and the Matrix Exponential

Consider a first order linear ordinary differential equations system in matrix form by

$$\frac{d\mathbf{x}(t)}{dt} = A\mathbf{x}(t), \mathbf{x}(0) = \mathbf{x}_0,$$

where $A \in \mathbb{R}^{n \times n}$ is a constant matrix and $\mathbf{x}(t) \in \mathbb{R}^n$ is the state vector.

The central mathematical object for solving such systems is the matrix exponential.

Definition 3.1 (Matrix Exponential)

For any square matrix $A \in \mathbb{R}^{n \times n}$, the matrix exponential is defined by the power series

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!}.$$

This convergence is ultimate to all square matrices so that the matrix exponential is well-defined no matter the size or norm of A.

The matrix exponential has a number of basic properties that are similar to those of the scalar exponential function.

Theorem 3.1.

Let $A, B \in \mathbb{R}^{n \times n}$. Then:

1. $e^0 = I$,
2. e^A is invertible with $(e^A)^{-1} = e^{-A}$,
3. If $AB = BA$, then $e^{A+B} = e^A e^B$.

Proof.

The results follow directly from the absolute convergence of the defining power series and standard properties of matrix multiplication.

A key analytical property of the matrix exponential is its behavior under differentiation.

Theorem 3.2.

For a constant matrix A ,

$$\frac{d}{dt}(e^{At}) = Ae^{At} = e^{At}A.$$

Proof.

Term-by-term differentiation of the power series defining e^{At} yields the result.

These properties establish the matrix exponential as a natural candidate for the fundamental solution operator of linear systems.

3.2 Fundamental Matrix, Norm Estimates, and Spectral Representation

A matrix-valued function $\Phi(t)$ is called a *fundamental matrix* of the linear system if it satisfies

$$\frac{d\Phi(t)}{dt} = A\Phi(t), \Phi(0) = I.$$

Theorem 3.3.

The fundamental matrix of the system $\dot{\mathbf{x}} = A\mathbf{x}$ is given by

$$\Phi(t) = e^{At}.$$

Proof.

Substituting $\Phi(t) = e^{At}$ into the defining conditions verifies both the differential equation and the initial condition.

Matrix norms are useful for analyzing stability and growth behavior of solutions.

Lemma 3.1.

For any consistent matrix norm $\|\cdot\|$,

$$\|e^A\| \leq e^{\|A\|}.$$

Proof.

The stated inequality can be obtained by using the sub multiplicativity of the matrix norm and the definition of the matrix exponential as a power series.

When the matrix A is a diagonalizable operation, there is a spectral decomposition of the matrix.

$$A = PDP^{-1},$$

where D is a diagonal matrix of eigenvalues. In this case, the matrix exponential can be expressed as

$$e^{At} = Pe^{Dt}P^{-1},$$

with e^{Dt} obtained by exponentiating each diagonal entry of D .

In the case that A is not diagonalizable, Jordan canonical forms are used, which leads to the presence of exponential functions multiplied by polynomial terms. These representations are useful in the qualitative behavior of the solutions, more specifically the stability and the long-term dynamics.

The findings here are used to define the theoretical framework, which is used to solve the exponential matrices. Those preliminaries form a direct basis to the explicit derivation of the solutions of linear systems and form the foundation of numerical methods like exponential time differencing and Krylov subspace approximations that were discussed in the following chapters.

4. LINEAR HOMOGENEOUS SYSTEMS OF DIFFERENTIAL EQUATIONS

In this part, the precise solution of linear homogeneous systems of ordinary differential equations is developed in terms of matrix exponentials. The findings herein provided represent the analytical heart of the matrix exponential based exponential solution methods and give a direct generalization of the scalar exponential solution of the first-order linear differential equations.

4.1 Exact Solution via the Matrix Exponential

Consider the linear homogeneous system

$$\frac{d\mathbf{x}(t)}{dt} = A\mathbf{x}(t), \mathbf{x}(0) = \mathbf{x}_0,$$

where $A \in \mathbb{R}^{n \times n}$ is a constant coefficient matrix.

Theorem 4.1 (Exact Solution of Linear Homogeneous Systems)

The unique solution of the above system is given by

$$\mathbf{x}(t) = e^{At} \mathbf{x}_0.$$

Proof.

Let $\mathbf{x}(t) = e^{At} \mathbf{x}_0$. Differentiating with respect to t yields

$$\frac{d\mathbf{x}(t)}{dt} = A e^{At} \mathbf{x}_0 = A \mathbf{x}(t).$$

Evaluating at $t = 0$ gives $\mathbf{x}(0) = I \mathbf{x}_0 = \mathbf{x}_0$.

Uniqueness follows from the Picard–Lindelöf theorem for linear systems.

Remark 4.1.

The matrix exponential acts as the evolution operator of the system, mapping the initial state $\mathbf{x}_0$ to the state at time t .

4.2 Stability and Qualitative Behavior

The qualitative behavior of solutions is governed by the spectral properties of the coefficient matrix A .

Theorem 4.2 (Stability Criterion).

In case the eigenvalues of A have exclusively negative real numbers, the trivial solution $\mathbf{x}(t) = \mathbf{0}$ is asymptotically stable.

Proof.

Exponential decay of the solution components in the case when the real parts of all the eigenvalues are negative has been observed in the spectral representation of the matrix exponential as t tends to infinity.

Remark 4.2.

Direct and explicit connection between system stability and eigenvalue distributions Matrix exponential formulations offer a clear and straightforward connection between the distribution of eigenvalues and system stability.

4.3 Diagonalizable and Non-Diagonalizable Systems

If the matrix A is diagonalizable, then

$$A = PDP^{-1},$$

where D is diagonal. The solution becomes

$$\mathbf{x}(t) = Pe^{Dt}P^{-1}\mathbf{x}_0,$$

where the exponential of D is obtained by exponentiating each diagonal entry.

In the case when A is not diagonalizable, we use Jordan canonical forms. The solution in this instance is that the factors of the polynomials multiply exponential terms which shows that the generalized eigenvectors are present.

Remark 4.3.

non-diagonalizable systems often exhibit transient growth even when asymptotically stable.

4.4 Illustrative Example

Example 4.1.

Consider the system

$$\frac{d}{dt}\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -2 & -3 \end{pmatrix}\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

The values of the eigen-coefficients of the coefficient matrix are -1 and -2. The two eigenvalues are negative and real; thus, the system is asymptotically stable. Matrix exponential is used to get the explicit solution.

$$\mathbf{x}(t) = e^{At}\mathbf{x}_0,$$

which describes exponential decay of both state variables.

Remark 4.4.

This example shows how the exponential building methods give analytical understanding of the dynamics of the system without necessarily having to rely on analytical step by step numerical integration.

4.5 Fundamental Matrix Interpretation

The solution operator e^{At} serves as the fundamental matrix of the system. For any two time instants t_1 and t_2 ,

$$\mathbf{x}(t_2) = e^{A(t_2-t_1)}\mathbf{x}(t_1),$$

highlighting the semigroup property of the solution.

Remark 4.5.

It is a critical property especially in the control theory and time-propagation algorithms.

5. NON-HOMOGENEOUS AND SEMI-LINEAR SYSTEMS

Most dynamical systems in practical situations are forced or otherwise influenced by nonlinear processes and thus become non-homogeneous or semi-linear systems of equations. Methods based on the matrix exponential can also be generalized to these systems and can be used to give a unified analysis framework to generalize the analysis of homogeneous linear systems.

Consider the non-homogeneous linear system

$$\frac{d\mathbf{x}(t)}{dt} = A\mathbf{x}(t) + \mathbf{f}(t), \mathbf{x}(0) = \mathbf{x}_0,$$

where $A \in \mathbb{R}^{n \times n}$ is a constant matrix and $\mathbf{f}(t) \in \mathbb{R}^n$ is a continuous forcing function. Using the variation-of-constants principle, the unique solution of this system can be expressed explicitly in terms of the matrix exponential as

$$\mathbf{x}(t) = e^{At}\mathbf{x}_0 + \int_0^t e^{A(t-s)}\mathbf{f}(s) ds.$$

The first term is the homogeneous response that is defined by the first condition and the integral term is the overall contribution of the external forcing, propagated locally by the system dynamics as the matrix exponential. In this context the matrix exponential can be considered as a Greens function of the linear system.

Semi-linear systems occur when there are nonlinear effects in presence of a linear operator and may be in the form.

$$\frac{d\mathbf{x}(t)}{dt} = A\mathbf{x}(t) + \mathbf{g}(\mathbf{x}(t)), \mathbf{x}(0) = \mathbf{x}_0,$$

where $\mathbf{g}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a nonlinear function. Applying the same variation-of-constants approach yields the exact integral representation

$$\mathbf{x}(t) = e^{At}\mathbf{x}_0 + \int_0^t e^{A(t-s)} \mathbf{g}(\mathbf{x}(s)) ds.$$

Though this is an exact expression, it is usually not calculable in closed form, because of the nonlinear term. However, it offers critical theory underlining the numerical techniques of matrix exponentials.

Specifically, time differencing techniques that are exponential in time are approximations to the nonlinear integral yet the linear term is treated exponentially exactly. The first-order exponential time differencing scheme is provided by.

$$\mathbf{x}_{n+1} = e^{Ah}\mathbf{x}_n + h \phi_1(Ah) \mathbf{g}(\mathbf{x}_n), \phi_1(z) = \frac{e^z - I}{z},$$

where h denotes the time step. Through this process of integrating the linear dynamics precisely, these types of schemes have superior stability properties than classical explicit methods particularly with the stiff systems.

When the eigenvalues of the matrix A are negative real values and the forcing or nonlinear term is bounded then the solution of the system is bounded in time. This is a stability property that underlines the strength

of the formulations based on matrix exponential and encourages their use in general in both analytical studies and numerical simulation of dynamical systems.

6. NUMERICAL APPROXIMATION VIA MATRIX EXPONENTIALS

Where the scale of the system or problem requires the evaluation of the matrix exponential to be computationally infeasible, then numerical approximation techniques are necessary. Numerical methods based on matrix exponential attempts to approximate the matrix exponential, or the action of the matrix exponential on vectors, and retain stability and accuracy properties of the exact solution.

The series expansion or diagonalization method of obtaining e^{At} is frequently not practical when there are many dimensions. Rather, the contemporary methods of numerical analysis are concerned with the calculations of expressions of the form $e^{At} \mathbf{v}$, where $\mathbf{v}$ is a specified vector. In this case, Krylov subspace techniques come in handy. Given a matrix A and a Vector $\mathbf{v}$, the Krylov subspace of dimension m is defined as

$$\mathcal{K}_m(A, \mathbf{v}) = \text{span}\{\mathbf{v}, A\mathbf{v}, A^2\mathbf{v}, \dots, A^{m-1}\mathbf{v}\}.$$

This low-dimensional subspace is then used to project the problem onto and obtain approximation of the matrix exponential with a large reduction of computational cost, without much loss of accuracy. Krylov convergence has been studied rigorously and error bounds have been provided to guarantee reliability in the case of large subspace dimensions.

The other significant numerical strategy is low-rank approximation of matrix differential equations. Solution matrices in many applications, e.g., control theory and quantum dynamics, have rapidly decaying singular values. Low-rank schemes take advantage of such a structure, by the solution being expressed as a lower-dimensional form and only the necessary components evolved. These methods, when used with matrix exponential methods, can be used to simulate otherwise intractable systems with high efficiency.

The major benefit of numerical schemes based on matrix exponential is stability. These methods are especially well suited to the stiff systems that are of interest to us as the linear component of the system is integrated without error, as opposed to the traditional explicit schemes which have particularly strict step-size limitations. This characteristic renders exponential integrators and Krylov based approximations

appealing to time-dependent problems that occur in physics, engineering and large-scale scientific computing.

7. APPLICATIONS AND DISCUSSION

Techniques based on matrices exponential are common to systems of differential equations analysis and simulation because of their capability to model system behavior precisely or structure preservingly. The use of matrix exponentials in linear control theory is associated with the analysis of state transition matrices, which allow straightforward analysis of the stability and system behavior according to spectral characteristics of the system matrix.

These techniques are especially useful in the fields of engineering and physical sciences in which stiff and high-dimensional systems are especially difficult to solve using classical numerical methods due to stability considerations. The combination of the linear component of the system is exact, leading to a better numerical robustness and accuracy of the matrix exponential based methods.

From a computational perspective, approaches like Krylov subspace methods and low rank formulations have greatly reduced the cost of computation of matrix exponentials. This has made it possible to use solvers based on exponentials in modern scientific computing environments.

Altogether, it is possible to state that the methods based on matrix exponential are characterized by a solid theoretical basis with practical computational benefits, which is why they are a stable and efficient approach to solving systems of differential equations in terms of diverse applications.

8. CONCLUSION

The methods based on matrix exponentials provide a rigorous and formulated way of solving sets of nonlinear equations of motion that simply connects the linear algebraic frameworks to the behavior of dynamical systems. The theoretical bases of matrix exponentials have been developed in a systematic way in this paper, and the extent of its usefulness to provide the exact solutions of the linear homogeneous systems has been extended to the non-homogeneous systems and semi-linear systems in terms of the integral representations. It was also indicated in the discussion that these analytical formulations support contemporary numerical methods, such as exponential time differencing, Krylov space approximations,

and low-rank methods, which can be efficiently and reliably computed to large-scale stiff systems. Combining classical theory and modern numerical techniques, the study highlights the versatility, stability and computational relevance of matrix exponential-based techniques to make them one of the key tools in the theoretical analysis and practical use of differential equations.

REFERENCES

1. Baleanu, D., Shiri, B., Srivastava, H. M., & Al Qurashi, M. (2018). *A Chebyshev spectral method based on operational matrix for fractional differential equations involving non-singular Mittag-Leffler kernel. Advances in Difference Equations*, 2018(1), 1-23.
2. Bloomfield, V. A. (2018). *Using R for numerical analysis in science and engineering. Chapman and Hall/CRC*.
3. Margossian, C. C. (2019). *A review of automatic differentiation and its efficient implementation. Wiley interdisciplinary reviews: data mining and knowledge discovery*, 9(4), e1305.
4. Mena, H., Ostermann, A., Pfurtscheller, L. M., & Piazzola, C. (2018). *Numerical low-rank approximation of matrix differential equations. Journal of Computational and Applied Mathematics*, 340, 602-614.
5. Subaşı, Y., Somma, R. D., & Orsucci, D. (2019). *Quantum algorithms for systems of linear equations inspired by adiabatic quantum computing. Physical review letters*, 122(6), 060504.
6. Yüzbaşı, Ş., & Sezer, M. (2013). *An exponential matrix method for solving systems of linear differential equations. Mathematical Methods in the Applied Sciences*, 36(3), 336-348.
7. Ma, W. X., Gu, X., & Gao, L. (2009). *A note on exact solutions to linear differential equations by the matrix exponential. Advances in Applied Mathematics and Mechanics*, 1(4), 573-580.
8. Al-Mohy, A. H., & Higham, N. J. (2011). *Computing the action of the matrix exponential, with an application to exponential integrators. SIAM journal on scientific computing*, 33(2), 488-511.
9. Çevik, M., Bahşı, M. M., & Sezer, M. (2014). *Solution of the delayed single degree of freedom system equation by exponential matrix method. Applied Mathematics and Computation*, 242, 444-453.
10. Sastre, J., Ibáñez, J., Defez, E., & Ruiz, P. (2011). *Accurate matrix exponential computation to solve coupled differential models in engineering. Mathematical and Computer Modelling*, 54(7-8), 1835-1840.

11. Al-Mohy, A. H., & Higham, N. J. (2010). *A new scaling and squaring algorithm for the matrix exponential. SIAM Journal on Matrix Analysis and Applications*, 31(3), 970-989.
12. Doha, E. H., Bhrawy, A. H., & Ezz-Eldien, S. S. (2011). *Efficient Chebyshev spectral methods for solving multi-term fractional orders differential equations. Applied Mathematical Modelling*, 35(12), 5662-5672.